

GAINS release notes

GAINS Developer Team, gains.dev@iiasa.ac.at

Release 4.01; September 2021

This is a major update of the model structure as well as elements of interface. It is a result of many strands of development that have taken place since the last version of GAINS that was made public in 2018. The principal features of the online model remain like before and the changes may not be obvious at first sight. Nonetheless, they will affect and, as we believe, improve the work of many users. This release addresses topics of growing importance such as the urban/rural distinction of activities and respective emissions from residential combustion and municipal waste management (including a much more detailed representation of the waste sector), non-compliant vehicles (“high emitters”) as well as explicit representation of international shipping and forest fires scenarios for use in impact assessment.

This is also the first release which is officially numbered. Internally, the existing previous interface version was known as version 3, therefore the numbering starts with 4.01. From now on, the versioning system will distinguish

- major updates which also change the design and functionality of the online model interface (i.e. all regional interfaces) as integer number corresponding to the online interface *version*,
- minor updates of functionality, bug fixes, etc., as *releases* within the same online model version, and might affect only individual regional interfaces.

Release 4.01 is the first release using online interfaces version 4.

Three categories of updates can be distinguished:

- Sectoral structure updated - valid for all regional interfaces. These are described in Section 1 below.
- Regional interface dependent updates are described in Section 2. These concerns mostly the impact calculations, changes in the region structure in GAINS-Europe, and map display/export features.
- Display features – see Section 2. These concern primarily higher spatial resolution of maps than before (0.1° outside Europe). The display can be switched between raster and smoothed contours, and every map can be exported in vector and GIS formats like GeoTIFF.

Available scenarios:

Release 4.01 includes a set of scenarios which have been recently published by IIASA's Pollution Management group (e.g., [EU Clean Air Outlook 2](#), [ECLIPSE V6b](#)) which can be used as a starting point for creating user defined scenarios.

Scenarios from previous versions:

Per default, **existing user defined scenarios are not ported** to GAINS version 4 due to the structural changes described below. The previous GAINS interfaces will remain active and accessible for the time being, but no further development is planned and only limited support can be provided by the IIASA team.

We understand that users may be interested to continue working with their existing scenarios in the new interface and make use of the enhanced capabilities of the new GAINS version. Depending on the age of the scenario (in particular, the emission vector used), this may be possible, but needs to be assessed on a case-by-case basis. Please get in touch with the developers' team (gains.dev@iiasa.ac.at) if you would like to have scenarios converted. We will try to provide assistance for a limited number of cases. Please note, however, that due to the structural changes the previous results will not be reproduced exactly in GAINS₄. Note that if your scenarios were developed based on ECLIPSE V4/5 in GAINS-3, we recommend you to directly create new scenarios in GAINS₄ based on the available default scenarios rather than convert your scenarios into GAINS₄, since ECLIPSE v4/5 are outdated.

1. Structural updates (interface independent)

| Topic | Previous implementation | Implementation in the current version of GAINS |
|---------------------------------|---|--|
| Model Release 4.01 | | |
| Municipal solid waste (MSW_XXX) | One sector: WASTE_RES (residential waste) | Multiple sectors specifying waste type and urban/rural residence: MSW_URB_[type] and MSW_RUR_[type] With [type] being: TEX: Textiles PAP: Paper PLA: Plastics WOOD: Wood FOOD: Food |

| | | |
|---------------------------------|---|---|
| | | <p>MET: Metals GLA: Glass OTH: Other</p> <p>Activity is specified in million tons of waste generated annually. Control options have been extended to include a wide range of different treatment of collected and uncollected waste. Calculations of air pollution and methane emissions have been harmonized.</p> |
| Residential combustion (DOM_xx) | <p>Activity: total DOM sector activity split into import sectors: DOM_RES (residential), DOM_COM (commercial), DOM_OTH (other).</p> <p>Control strategy and emission factors are specified at the level of sectors like: DOM_STOVE_C, DOM_STOVE_H,...</p> <p>Sub-structure for activity splits into cooking and heating stoves for solid fuels: DOM_RES_STOVE_C, DOM_RES_STOVE_H, DOM_COM_STOVE_C, DOM_COM_STOVE_H, DOM_OTH_STOVE_C, DOM_OTH_STOVE_H</p> | <p>Activity: Urban-rural split introduced in residential combustion: total DOM activity is distributed into import sectors DOM_URB, DOM_RUR, DOM_COM, DOM_OTH (Note: DOM_RES=DOM_URB+DOM_RUR does not exist as a separate sector anymore). By default, “urban” and “rural” correspond to UN World Urbanization Prospects 2018 (downscaled to GAINS regions).</p> <p>Control strategy and emission factors are specified at the level of sectors as before: DOM_STOVE_C, DOM_STOVE_H,...</p> <p>Sub-structure for activity splits into cooking (C) and heating (H) stoves for solid fuels: DOM_URB_STOVE_C, DOM_URB_STOVE_H, DOM_RUR_STOVE_C, DOM_RUR_STOVE_H, DOM_COM_STOVE_C, DOM_COM_STOVE_H, DOM_OTH_STOVE_C, DOM_OTH_STOVE_H</p> |
| High emitting vehicles | Not included in the online model. Included in gridded global emission products for the atmospheric modelling community, such as the ECLIPSE data sets. | <p>Included in the online model as an approximation to the calculations done for the gridded emission products.</p> <p>New activity classes: MD_NV_HE: non-complying diesel vehicles GSL_NV_HE: non-complying gasoline vehicles</p> |

| | | |
|------------------------|--|---|
| | | The activity is derived from total numbers of vehicles driving with these fuels (MD_NV, GSL_NV) through shares of non-compliant vehicles. Corresponding emission factors are calculated in the emission vector initialization as average over the control technologies applied in a selected representative scenario. |
| International shipping | <p>In GAINS-Europe domain: Emission-only scenarios for 11 sea regions.</p> <p>Outside GAINS-Europe: not included in the online model; included in gridded global emission products for the atmospheric modelling community, such as the ECLIPSE data sets.</p> | <p>In GAINS-Europe domain: Emission-only scenarios for 13 sea regions as described in Cofala et al. (2018)</p> <p>Outside GAINS-Europe: We included International Shipping as a new element in the scenario meta data (previously “Scenario description”) affecting only ambient PM_{2.5} concentrations outside the European domain. Different pre-set storylines can be chosen. A default is allocated to any new scenario.</p> |
| Forest fires | Included in the natural contributions to ambient PM2.5, constant over time. | Included in the natural contributions to ambient PM2.5. Variable over time, following pre-set storylines which can be selected in the scenario metadata (previously “Scenario description”). A default storyline is allocated to any new scenario. |
| Cost calculation | All costs specified in Euros 2005. Fixed storyline for wages and energy prices. | All costs specified in Euros 2015. A storyline on wages and fuel prices can be selected in the scenario metadata (previously “Scenario description”). Available storylines are: .) dynamic prices for 2025 .) constant prices based on WEO 2009 .) dynamic prices based on WEO 2009 |

2. Interface specific changes

| Topic | Interface version | Previous implementation | Implementation in current version of GAINS |
|---------------------------|---------------------|---|--|
| Model release 4.01 | | | |
| Model regions | Europe | <ul style="list-style-type: none"> • 41 regions • Serbia, Montenegro and Kosovo combined as one region SEMO_WHOL. | <ul style="list-style-type: none"> • 45 regions • SEMO_WHOL split into Serbia (SERB_WHOL), Montenegro (MONT_WHOL), Kosovo (KOSO_WHOL) • Georgia (GEOR_WHOL), Armenia (ARME_WHOL), Azerbaijan (AZER_WHOL) added as model regions |
| | Global (not public) | <ul style="list-style-type: none"> • 174 regions • United States: one region (USAM_WHOL) • Africa: only Egypt (EGYP_WHOL), Northern Africa (NAFR_WHOL), South Africa (SAFR_WHOL), and Other Africa (OAFR_WHOL) | <ul style="list-style-type: none"> • 180 regions, including those in the public interfaces for Europe (changes see above) and Asia. • United States have been split into Mainland (including Hawaii) USAM_MAIN and Alaska USAM_ALAS • Africa has been split into nine regions: Egypt (EGYP_WHOL), South Africa (SAFR_WHOL), Nigeria (NIGE_WHOL), Tanzania (TANZ_WHOL), Kenya (KENY_WHOL), Northern Africa (NAFR_WHOL), Western Africa (WAFR_WHOL), Eastern Africa (EAFR_WHOL), Remaining Southern Africa (RSAF_WHOL). Sudan has been moved from NAFR_WHOL to EAFR_WHOL. |

| | | | |
|------------------------------------|-----------------------|--|--|
| Ambient PM and impact calculations | Europe | <ul style="list-style-type: none"> • Atmospheric transfer coefficients based on EMEP CTM (~28km resolution) with downscaling to 7km based on CHIMERE CTM. • Life expectancy loss calculated based on life table of UN World Population Prospects 2010 • Premature deaths calculated for 2010 population | <ul style="list-style-type: none"> • Transfer coefficients (same methodology) extended to include the new regions and updated sea regions. • Life table used for life expectancy loss can be switched between UN World Population Prospects 2010 and 2017 editions. • Premature deaths can be calculated for 2010 population (“static”) and scenario year dependent population. |
| | East Asia, South Asia | <ul style="list-style-type: none"> • Atmospheric transfer coefficients based on TM5 CTM (1° resolution) • Life expectancy loss calculated using linear dose-response function | <ul style="list-style-type: none"> • Atmospheric transfer coefficients based on EMEP CTM (0.5° resolution / 0.1° for low-level sources) • Premature deaths attributable to ambient PM2.5 calculated using integrated exposure-response relationships (IERs) from Global Burden of Disease 2013 / WHO (2016) |
| | Asia | <ul style="list-style-type: none"> • Atmospheric transfer coefficients based on EMEP CTM (0.5° resolution), downscaling to urban polygons for low-level primary PM emission sources. • Ambient PM2.5 maps at 0.5° resolution. • No health impact calculations. | <ul style="list-style-type: none"> • Atmospheric transfer coefficients based on EMEP CTM (0.5° resolution / 0.1° for low-level sources) • Ambient PM2.5 maps at 0.1° resolution. • Premature deaths attributable to ambient PM2.5 calculated using integrated exposure-response relationships (IERs) from Global Burden of Disease 2013 / WHO (2016). |